

Microarray Analysis Reveals Distinct Gene Expression Profiles among Different Histologic Types of Endometrial Cancer

John I. Risinger, G. Larry Maxwell, G. V. R. Chandramouli, Amir Jazaeri, Olga Aprelikova, Tricia Patterson, Andrew Berchuck, and J. Carl Barrett*

*Laboratory of Biosystems and Cancer, National Cancer Institute, Bethesda, Maryland 20892 [J. I. R., G. L. M., G. V. R. C., A. J., O. A., T. P., J. C. B.];
Walter Reed Army Medical Center, Washington, DC 20013 [G. L. M];
Department of Obstetrics and Gynecology/Division of Gynecologic Oncology, Duke University, Durham, North Carolina 27710 [A. B.]*

CANCER RESEARCH, **63**, 6–11, January 1, 2003

Expression data description

Expressions ratios are given relative to geometric average of 7 normal endometrial samples. Gene identifiers were obtained from mAdb (microarray database at National Cancer Institute, NIH, <http://nciarray.nci.nih.gov>). All the identifiers except “Unique Id” are periodically updated by mAdb to maintain accurate book-keeping up to date.

- Unique Id: A unique identifier assigned to each spot on microarray by mAdb to track the probe.
- Clone Id: Incyte PD clone identification number.
- UniGene cluster: NCBI UniGene cluster number.
- Gene symbol: Official gene symbol.
- Description: The gene description corresponding to BLAST.